

Recombinant Human CD3D cell lysate

Cat. No. CD3D-1274HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CD3D / CD3-Delta derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	CD3D CD3d molecule, delta (CD3-TCR complex) [Homo sapiens]
Official Symbol	CD3D
Synonyms	CD3D; CD3d molecule, delta (CD3-TCR complex); CD3d antigen, delta polypeptide (TiT3 complex) , T3D; T-cell surface glycoprotein CD3 delta chain; CD3 delta; OKT3, delta chain; CD3 antigen, delta subunit; T-cell receptor T3 delta chain; CD3d antigen, delta polypeptide (TiT3 complex); T3D; CD3-DELTA;
Gene ID	915
mRNA Refseq	NM_000732
Protein Refseq	NP_000723
MIM	186790
UniProt ID	P04234

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	11q23
Pathway	Adaptive Immune System, organism-specific biosystem; CXCR4-mediated signaling events, organism-specific biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Costimulation by the CD28 family, organism-specific biosystem; Downstream TCR signaling, organism-specific biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem;
Function	protein heterodimerization activity; receptor activity; transcription coactivator activity; transmembrane signaling receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA