Recombinant Human PDIA4 cell lysate

Cat. No. PDIA4-1285HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human ERP72 / PDIA4 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	PDIA4 protein disulfide isomerase family A, member 4 [Homo sapiens]
Official Symbol	PDIA4
Synonyms	PDIA4; protein disulfide isomerase family A, member 4; protein disulfide isomerase associated 4; protein disulfide-isomerase A4; ERP70; ERP72; ER protein 70; ER protein 72; protein disulfide isomerase-associated 4; endoplasmic reticulum resident protein 70; endoplasmic reticulum resident protein 72; protein disulfide isomerase related protein (calcium-binding protein, intestinal-related); ERp-72;
Gene ID	9601
mRNA Refseq	NM_004911
Protein Refseq	NP_004902
UniProt ID	P13667

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	7q35
Pathway	Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem; Vibrio cholerae infection, organism-specific biosystem; Vibrio cholerae infection, conserved biosystem;
Function	electron carrier activity; isomerase activity; protein disulfide isomerase activity; protein disulfide oxidoreductase activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA