

Recombinant Human B3GNT2 cell lysate

Cat. No. B3GNT2-1396HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human B3GNT2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

B3GNT2 UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 2 [Homo sapiens]

Official Symbol

B3GNT2

Synonyms

B3GNT2; UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 2; B3GNT1, UDP GlcNAc:betaGal beta 1,3 N acetylglucosaminyltransferase 1; B3GN T1; B3GN T2; B3GNT 2; BETA3GNT; beta3Gn-T1; beta3Gn-T2; beta-1,3-Gn-T1; beta-1,3-Gn-T2; beta-1,3-N-acetylglucosaminyltransferase bGnT-1; beta-1,3-N-acetylglucosaminyltransferase bGnT-2; UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 1; B3GNT; BGNT2; B3GNT1; BGnT-2; B3GN-T2; B3GNT-2;

Gene ID

10678

mRNA Refseq

NM_006577

Protein Refseq

NP_006568

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	605581
UniProt ID	Q9NY97
Chromosome Location	2p15
Pathway	Glycosaminoglycan biosynthesis - keratan sulfate, organism-specific biosystem; Glycosaminoglycan biosynthesis - keratan sulfate, conserved biosystem; Glycosphingolipid biosynthesis - lacto and neolacto series, organism-specific biosystem; Glycosphingolipid biosynthesis - lacto and neolacto series, conserved biosystem; Metabolic pathways, organism-specific biosystem; Metabolism of proteins, organism-specific biosystem; O-linked glycosylation of mucins, organism-specific biosystem;
Function	UDP-galactose:beta-N-acetylglucosamine beta-1,3-galactosyltransferase activity; galactosyltransferase activity; transferase activity, transferring glycosyl groups;