Recombinant Human AGER cell lysate

Cat. No. AGER-1480HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human AGER / RAGE derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	AGER advanced glycosylation end product-specific receptor [Homo sapiens]
Official Symbol	AGER
Synonyms	AGER; advanced glycosylation end product-specific receptor; RAGE; RAGE isoform sRAGE-delta; RAGE isoform NtRAGE-delta;
Gene ID	177
mRNA Refseq	NM_001136
Protein Refseq	NP_001127
MIM	600214
UniProt ID	Q15109
Chromosome Location	6p21.3

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Activated TLR4 signalling, organism-specific biosystem; Advanced glycosylation endproduct receptor signaling, organism-specific biosystem; Cytosolic sensors of pathogen-associated DNA, organism-specific biosystem; DAI mediated induction of type I IFNs, organism-specific biosystem; Immune System, organism-specific biosystem; Innate Immune System, organism-specific biosystem; MyD88 cascade initiated on plasma membrane, organism-specific biosystem;
Function	S100 alpha binding; advanced glycation end-product receptor activity; protein binding; receptor activity; receptor activity; transmembrane signaling receptor activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA