

Recombinant Human ENPP2 cell lysate

Cat. No. ENPP2-1556HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human Autotaxin / ENPP2 / NPP2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

ENPP2 ectonucleotide pyrophosphatase/phosphodiesterase 2 [Homo sapiens]

Official Symbol

ENPP2

Synonyms

ENPP2; ectonucleotide pyrophosphatase/phosphodiesterase 2; PDNP2; ectonucleotide pyrophosphatase/phosphodiesterase family member 2; ATX; autotaxin; PD IALPHA; E-NPP 2; autotaxin-t; plasma lysophospholipase D; extracellular lysophospholipase D; phosphodiesterase I/nucleotide pyrophosphatase 2; NPP2; ATX-X; LysoPLD; AUTOTAXIN; PD-IALPHA; FLJ26803;

Gene ID

5168

mRNA Refseq

NM_001040092

Protein Refseq

NP_001035181

MIM

601060

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q13822
Chromosome Location	8q24.1
Pathway	Ether lipid metabolism, organism-specific biosystem; Ether lipid metabolism, conserved biosystem;
Function	alkylglycerophosphoethanolamine phosphodiesterase activity; calcium ion binding; hydrolase activity; lysophospholipase activity; nucleic acid binding; nucleotide diphosphatase activity; phosphodiesterase I activity; polysaccharide binding; scavenger receptor activity; transcription factor binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA