

Recombinant Human BLNK cell lysate

Cat. No. BLNK-1858HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human BLNK derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	BLNK B-cell linker [Homo sapiens]
Official Symbol	BLNK
Synonyms	BLNK; B-cell linker; B-cell linker protein; B cell adaptor containing SH2 domain; B cell activation; B cell adapter containing a SH2 domain protein; BASH; bca; BLNK s; Ly57; SLP 65; SLP65; Src homology [SH2] domain containing leukocyte protein of 65 kD; B-cell activation; B cell linker protein; cytoplasmic adapter protein; B-cell adapter containing a SH2 domain protein; B-cell adapter containing a Src homology 2 domain protein; Src homology 2 domain-containing leukocyte protein of 65 kDa; Src homology [SH2] domain-containing leukocyte protein of 65 kD; AGM4; LY57; BLNK-S; SLP-65; MGC111051;
Gene ID	29760
mRNA Refseq	NM_001114094
Protein Refseq	NP_001107566

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	604515
UniProt ID	Q8WV28
Chromosome Location	10q23.2-q23.33
Pathway	Adaptive Immune System, organism-specific biosystem; Antigen Activates B Cell Receptor Leading to Generation of Second Messengers, organism-specific biosystem; B Cell Receptor Signaling Pathway, organism-specific biosystem; B cell receptor signaling pathway, organism-specific biosystem; B cell receptor signaling pathway, conserved biosystem; BCR signaling pathway, organism-specific biosystem; Cytokine Signaling in Immune system, organism-specific biosystem;
Function	SH3/SH2 adaptor activity; SH3/SH2 adaptor activity; protein binding; transmembrane receptor protein tyrosine kinase adaptor activity;