Active Recombinant Human Chemokine (C-C Motif) Ligand 21, MIgG2a Fc-tagged

Cat. No. CCL21-122H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CCL21 (AAH27918.1)(Ser24-Pro134) is fused to the N-terminus of the Fc region of mouse IgG2a was expressed in CHO cell.
Species	Human
Source	CHO
ProteinLength	24-134 a.a.
Description	This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. Similar to other chemokines the protein encoded by this gene inhibits hemopoiesis and stimulates chemotaxis. This protein is chemotactic in vitro for thymocytes and activated T cells, but not for B cells, macrophages, or neutrophils. The cytokine encoded by this gene may also play a role in mediating homing of lymphocytes to secondary lymphoid organs. It is a high affinity functional ligand for chemokine receptor 7 that is expressed on T and B lymphocytes and a known receptor for another member of the cytokine family (small inducible cytokine A19)
Form	Lyophilized from 0.2μm-filtered solution in PBS.
Bio-activity	Measured by its ability to chemoattract 5-10 day cultured human peripheral blood

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	lymphocytes (PBL).
Molecular Mass	38KDa (monomer)
AA Sequence	Ser24-Pro134
Endotoxin	<0.06 eu/μg="" as="" determined="" by="" lal="">
Purity	>98%, by SDS-PAGE under reducing conditions.
Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Reconstitution	Reconstitute at 100μg/ml in sterile PBS.
Warning	Avoid freeze/thaw cycles.

GENE INFORMATION

Gene Name	CCL21 chemokine (C-C motif) ligand 21 [Homo sapiens]
Official Symbol	CCL21
Synonyms	CCL21; chemokine (C-C motif) ligand 21; SCYA21, small inducible cytokine subfamily A (Cys Cys), member 21; C-C motif chemokine 21; 6Ckine; beta chemokine exodus 2; CKb9; ECL; Efficient Chemoattractant for Lymphocytes; exodus 2; secondary lymphoid tissue chemokine; SLC; TCA4; exodus-2; beta chemokine exodus-2; beta-chemokine exodus-2; small-inducible cytokine A21; secondary lymphoid-tissue chemokine; small inducible cytokine subfamily A (Cys-Cys), member 21; SCYA21; MGC34555;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	6366
mRNA Refseq	NM_002989
Protein Refseq	NP_002980
MIM	602737
UniProt ID	O00585
Chromosome Location	9p13
Pathway	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (i) signalling events, organism-specific biosystem;
Function	CCR7 chemokine receptor binding; chemokine activity; chemokine activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA