

Recombinant Human Chemokine (C-C Motif) Ligand 2

Cat. No. CCL2-124H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Mammalian Cells
Description	Monocyte chemotactic protein-1 (MCP-1) is an essential chemokine produced predominantly by macrophages and endothelial cells and is a potent chemotactic factor for monocytes. It can induce the proliferation and activation of killer cells known as CHAK (CC-Chemokine-activated killer). MCP-1 is also a potent activator of human basophils, induces degranulation and the release of histamines. MCP-1 is involved in the pathogenesis of atherosclerosis and myocardial infarction.
State Of Matter	Lyophilized.
Purity	>97% by SDS Page and analyzed by silver stain.
Endotoxin	<0.1 ng per 1 g as determined by the LAL method.
Molecular Weight	The predicted molecular weight of Recombinant Human MCP-1 is Mr 35 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 70-95 kDa.
Storage And Stability	This lyophilized protein is stable for six to twelve months when stored desiccated at -20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

for one month or at -20°C to -70°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

GENE INFORMATION

Gene Name CCL2 chemokine (C-C motif) ligand 2 [Homo sapiens]

Synonyms

CCL2; chemokine (C-C motif) ligand 2; GDCF-2; HC11; HSMCR30; MCAF; MCP-1; MCP1; MGC9434; SCYA2; SMC-CF; monocyte chemoattractant protein-1; monocyte chemotactic and activating factor; monocyte secretory protein JE; small inducible cytokine A2; small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig-je); small inducible cytokine subfamily A (Cys-Cys), member 2; Monocyte chemoattractant protein 1; Monocyte chemotactic and activating factor; Monocyte chemotactic protein 1; Monocyte secretory protein JE; Small-inducible cytokine A2

Gene ID 6347

mRNA Refseq NM_002982

Protein Refseq NP_002973

MIM 158105

UniProt ID P13500

Chromosome Location 17q11.2-q12

Pathway Cytokine-cytokine receptor interaction; Chemokine signaling pathway

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Function

CCR2 chemokine receptor binding; G-protein-coupled receptor binding; chemokine activity; heparin binding; signal transducer activity; protein kinase activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA