

Recombinant Human Chemokine (C-C Motif) Ligand 5

Cat. No. CCL5-125H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Mammalian Cells
Description	RANTES also known as Chemokine (C-C motif) ligand 5 (CCL5), a CC-chemokine is an important mediators of the immune and inflammatory response. It is also a potent chemoattractant for T cells and eosinophils. RANTES is involved in many inflammatory diseases, for example, bronchial asthma, delayed-type hypersensitivity reactions, viral infections, arthritis, chronic eosinophilic pneumonia, and idiopathic interstitial pneumonia.
State Of Matter	Lyophilized.
Purity	>90% by SDS Page and analyzed by silver stain.
Endotoxin	<1.0 EU/g as determined by the LAL method.
Molecular Weight	The predicted molecular weight of Recombinant Human RANTES is Mr 34 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 85-105 kDa.
Storage And Stability	This lyophilized protein is stable for six to twelve months when stored desiccated at -20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

for one month or at -20°C to -70°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

GENE INFORMATION

Gene Name CCL5 chemokine (C-C motif) ligand 5 [Homo sapiens]

Synonyms

CCL5; chemokine (C-C motif) ligand 5; SISd; beta-chemokine RANTES; EoCP; RANTES; TCP228; MGC17164; SCYA5; RANTES; TCP228; D17S136E; MGC17164; chemokine (C-C motif) ligand 5; regulated upon activation; normally T-expressed; and presumably secrete; presumably secreted; small inducible cytokine A5; small inducible cytokine A5 (RANTES); small inducible cytokine subfamily A (Cys-Cys); member 5; Small-inducible cytokine A5; Small-inducible cytokine A5; T cell-specific protein P228; Eosinophil-chemotactic cytokine; RANTES(3-68); RANTES(4-68); T-cell specific RANTES protein; T-cell-specific protein RANTES; SIS-delta

Gene ID 6352

mRNA Refseq NM_002985

Protein Refseq NP_002976

MIM 187011

UniProt ID P13501

Chromosome Location 17q11.2-q12

Pathway

Chemokine signaling pathway; Cytokine-cytokine receptor interaction; Epithelial cell signaling in Helicobacter pylori infection; Prion diseases; Toll-like receptor signaling

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

pathway; Signaling by GPCR

Function

chemoattractant activity; chemokine activity; signal transducer activity

PDB rendering based
on 1pmi.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA