

Recombinant Human FGFR4 cell lysate

Cat. No. FGFR4-1907HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human FGFR4 / FGF Receptor 4 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

FGFR4 fibroblast growth factor receptor 4 [Homo sapiens]

Official Symbol

FGFR4

Synonyms

FGFR4; fibroblast growth factor receptor 4; CD334; JTK2; FGFR-4; tyrosylprotein kinase; protein-tyrosine kinase; hydroxyaryl-protein kinase; tyrosine kinase related to fibroblast growth factor receptor; TKF; MGC20292;

Gene ID

2264

mRNA Refseq

NM_002011

Protein Refseq

NP_002002

MIM

134935

UniProt ID

P22455

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	5q33-qter
Pathway	Downstream signaling of activated FGFR, organism-specific biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved biosystem; FGF signaling pathway, organism-specific biosystem; FGFR ligand binding and activation, organism-specific biosystem; FGFR4 ligand binding and activation, organism-specific biosystem; FRS2-mediated cascade, organism-specific biosystem;
Function	ATP binding; fibroblast growth factor 1 binding; fibroblast growth factor 2 binding; fibroblast growth factor binding; fibroblast growth factor-activated receptor activity; fibroblast growth factor-activated receptor activity; heparin binding; nucleotide binding; protein tyrosine kinase activity; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA