

Recombinant Human MST1R cell lysate

Cat. No. MST1R-1962HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human MST1R / CD136 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	MST1R macrophage stimulating 1 receptor (c-met-related tyrosine kinase) [Homo sapiens]
Official Symbol	MST1R
Synonyms	MST1R; macrophage stimulating 1 receptor (c-met-related tyrosine kinase); PTK8, PTK8 protein tyrosine kinase 8 , RON; macrophage-stimulating protein receptor; CD136; CDw136; p185-Ron; MSP receptor; RON variant E2E3; MST1R variant RON30; MST1R variant RON62; soluble RON variant 1; soluble RON variant 2; soluble RON variant 3; soluble RON variant 4; c-met-related tyrosine kinase; PTK8 protein tyrosine kinase 8; RON; PTK8;
Gene ID	4486
mRNA Refseq	NM_001244937
Protein Refseq	NP_001231866

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	600168
UniProt ID	Q04912
Chromosome Location	3p21
Pathway	Alpha6-Beta4 Integrin Signaling Pathway, organism-specific biosystem; a6b1 and a6b4 Integrin signaling, organism-specific biosystem; amb2 Integrin signaling, organism-specific biosystem;
Function	ATP binding; enzyme binding; macrophage colony-stimulating factor receptor activity; nucleotide binding; protein binding; receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA