Recombinant Rat FCGR1 cell lysate

Cat. No. FCGR1-1972RCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Rat CD64 / FCGR1A derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Rat
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Fcgr1a Fc fragment of IgG, high affinity Ia, receptor (CD64) [Rattus norvegicus]
Official Symbol	Fcgr1a
Synonyms	Fcgr1; Fcgr1b; FcgammaRI; high affinity immunoglobulin gamma Fc receptor I; Fc receptor, IgG, high affinity I; high affinity IgG Fc receptor, subunit beta; Fc fragment of IgG high affinity Ib receptor; high-affinity immunoglobulin gamma Fc receptor I; Fc fragment of IgG, high affinity Ib, receptor for (CD64)
Gene ID	295279
mRNA Refseq	NM_001100836
Protein Refseq	NP_001094306
Chromosome Location	2q34

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Adaptive Immune System, organism-specific biosystem; Fc gamma R-mediated phagocytosis, conserved biosystem; Innate Immune System, organism-specific biosystem
Function	IgG binding; IgG receptor activity

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA