

Recombinant Human HMGB1 cell lysate

Cat. No. HMGB1-2076HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human HMGB1 / HMG1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	HMGB1 high mobility group box 1 [Homo sapiens]
Official Symbol	HMGB1
Synonyms	HMGB1; high mobility group box 1; high mobility group (nonhistone chromosomal) protein 1 , high mobility group box 1 , HMG1; high mobility group protein B1; Amphoterin; DKFZp686A04236; high mobility group protein 1; HMG3; SBP 1; Sulfoglucuronyl carbohydrate binding protein; HMG-1; high-mobility group box 1; high-mobility group (nonhistone chromosomal) protein 1; HMG1; SBP-1;
Gene ID	3146
mRNA Refseq	NM_002128
Protein Refseq	NP_002119
MIM	163905

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P09429
Chromosome Location	13q12
Pathway	Activated TLR4 signalling, organism-specific biosystem; Activation of DNA fragmentation factor, organism-specific biosystem; Advanced glycosylation endproduct receptor signaling, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis induced DNA fragmentation, organism-specific biosystem; Apoptotic executionphase, organism-specific biosystem;
Function	DNA binding; DNA binding, bending; DNA binding, bending; RAGE receptor binding; calcium-dependent protein kinase regulator activity; chemoattractant activity; cytokine activity; cytokine activity; damaged DNA binding; double-stranded DNA binding; protein binding; protein kinase activator activity; repressing transcription factor binding; sequence-specific DNA binding transcription factor activity; single-stranded DNA binding; transcription factor binding;