Recombinant Human SERPINI2 cell lysate

Cat. No. SERPINI2-2083HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human SERPINI2 / SerpinI2 / MEPI derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	SERPINI2 serpin peptidase inhibitor, clade I (pancpin), member 2 [Homo sapiens]
Official Symbol	SERPINI2
Synonyms	SERPINI2; serpin peptidase inhibitor, clade I (pancpin), member 2; PI14, serine (or cysteine) proteinase inhibitor, clade I (neuroserpin), member 2 , serine (or cysteine) proteinase inhibitor, clade I (pancpin), member 2; serpin I2; MEPI; PANCPIN; pancpin; TSA2004; PI-14; protease inhibitor 14; peptidase inhibitor 14; pancreas-specific protein TSA2004; myoepithelium-derived serine protease inhibitor; serine (or cysteine) proteinase inhibitor, clade I (pancpin), member 2; serine (or cysteine) proteinase inhibitor, clade I (neuroserpin), member 2; PI14;
Gene ID	5276
mRNA Refseq	NM_006217
Protein Refseq	NP_006208

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	605587
UniProt ID	O75830
Chromosome Location	3q26
Function	peptidase inhibitor activity; serine-type endopeptidase inhibitor activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA