Recombinant Mouse P4HB cell lysate

Cat. No. P4HB-2121MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse P4HB / ERBA2L derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	P4hb prolyl 4-hydroxylase, beta polypeptide [Mus musculus]
Official Symbol	P4HB
Synonyms	P4HB; prolyl 4-hydroxylase, beta polypeptide; protein disulfide-isomerase; p55; ER protein 59; protein disulfide isomerase; prolyl 4-hydroxylase subunit beta; cellular thyroid hormone-binding protein; endoplasmic reticulum resident protein 59; PDI; Thbp; ERp59; Pdia1;
Gene ID	18453
mRNA Refseq	NM_011032
Protein Refseq	NP_035162
Pathway	Chylomicron-mediated lipid transport, organism-specific biosystem; Lipid digestion, mobilization, and transport, organism-specific biosystem; Lipoprotein metabolism, organism-specific biosystem; Metabolism, organism-specific biosystem; Metabolism

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	of lipids and lipoproteins, organism-specific biosystem; Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem;
Function	electron carrier activity; isomerase activity; procollagen-proline 4-dioxygenase activity; protein disulfide isomerase activity; protein disulfide oxidoreductase activity; protein heterodimerization activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA