

Recombinant Human CES2 cell lysate

Cat. No. CES2-2139HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CES2 / Carboxylesterase-2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	CES2 carboxylesterase 2 [Homo sapiens]
Official Symbol	CES2
Synonyms	CES2; carboxylesterase 2; carboxylesterase 2 (intestine, liver); cocaine esterase; CE 2; CES2A1; iCE; hCE-2; methylumbelliferyl-acetate deacetylase 2; intestinal carboxylesterase; liver carboxylesterase-2; CE-2; PCE-2;
Gene ID	8824
mRNA Refseq	NM_003869
Protein Refseq	NP_003860
MIM	605278
UniProt ID	O00748

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	16q22.1
Pathway	Drug metabolism - other enzymes, organism-specific biosystem; Drug metabolism - other enzymes, conserved biosystem; E2F transcription factor network, organism-specific biosystem; Fluoropyrimidine Activity, organism-specific biosystem; Irinotecan Pathway, organism-specific biosystem;
Function	carboxylesterase activity; hydrolase activity; methyl indole-3-acetate esterase activity; methyl jasmonate esterase activity; methyl salicylate esterase activity; methylumbelliferyl-acetate deacetylase activity; retinyl-palmitate esterase activity;