

Recombinant Human TNFRSF10A cell lysate

Cat. No. TNFRSF10A-2180HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human TNFRSF10A / CD261 / APO2 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	TNFRSF10A tumor necrosis factor receptor superfamily, member 10a [Homo sapiens]
Official Symbol	TNFRSF10A
Synonyms	TNFRSF10A; tumor necrosis factor receptor superfamily, member 10a; tumor necrosis factor receptor superfamily member 10A; Apo2; CD261; DR4; TRAILR 1; TRAIL-R1; TRAIL receptor 1; death receptor 4; cytotoxic TRAIL receptor; TNF-related apoptosis-inducing ligand receptor 1; tumor necrosis factor receptor superfamily member 10a variant 2; APO2; TRAILR1; TRAILR-1; MGC9365;
Gene ID	8797
mRNA Refseq	NM_003844
Protein Refseq	NP_003835
MIM	603611

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	O00220
Chromosome Location	8p21
Pathway	Apoptosis, organism-specific biosystem; Apoptosis, conserved biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Direct p53 effectors, organism-specific biosystem; Influenza A, organism-specific biosystem; Influenza A, conserved biosystem;
Function	TRAIL binding; death receptor activity; protein binding; receptor activity; transcription factor binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA