Recombinant Human HYOU1 cell lysate

Cat. No. HYOU1-2295HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human HYOU1 / HSP12A derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	HYOU1 hypoxia up-regulated 1 [Homo sapiens]
Official Symbol	HYOU1
Synonyms	HYOU1; hypoxia up-regulated 1; hypoxia up-regulated protein 1; glucose regulated protein 170; Grp170; HSP12A; ORP150; glucose-regulated protein 170; 150 kDa oxygen-regulated protein; oxygen regulated protein (150kD); 170 kDa glucose-regulated protein; GRP-170; ORP-150; FLJ94899; FLJ97572; DKFZp686N08236;
Gene ID	10525
mRNA Refseq	NM_001130991
Protein Refseq	NP_001124463
MIM	601746
UniProt ID	Q9Y4L1

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	11q23.1-q23.3
Pathway	Activation of Chaperone Genes by XBP1(S), organism-specific biosystem; Activation of Chaperones by IRE1alpha, organism-specific biosystem; Diabetes pathways, organism-specific biosystem; Disease, organism-specific biosystem; Protein processing in endoplasmic reticulum, organism-specific biosystem; Protein processing in endoplasmic reticulum, conserved biosystem; Unfolded Protein Response, organism-specific biosystem;
Function	ATP binding; nucleotide binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA