

Recombinant Human FZD5 cell lysate

Cat. No. FZD5-2388HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human Frizzled-5 / FZD5 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[FZD5 frizzled family receptor 5 \[Homo sapiens \]](#)

Official Symbol

FZD5

Synonyms

FZD5; frizzled family receptor 5; C2orf31, chromosome 2 open reading frame 31 , frizzled (Drosophila) homolog 5 , frizzled 5, seven transmembrane spanning receptor , frizzled homolog 5 (Drosophila); frizzled-5; DKFZP434E2135; HFZ5; fz-5; fzE5; Wnt receptor; seven-transmembrane receptor frizzled-5; frizzled 5, seven transmembrane spanning receptor; C2orf31; MGC129692; DKFZp434E2135;

Gene ID

[7855](#)

mRNA Refseq

[NM_003468](#)

Protein Refseq

[NP_003459](#)

MIM

[601723](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q13467
Chromosome Location	2q33.3
Pathway	Basal cell carcinoma, organism-specific biosystem; Basal cell carcinoma, conserved biosystem; Canonical Wnt signaling pathway, organism-specific biosystem; Class B/2 (Secretin family receptors), organism-specific biosystem; GPCR ligand binding, organism-specific biosystem; HTLV-I infection, organism-specific biosystem; HTLV-I infection, conserved biosystem;
Function	G-protein coupled receptor activity; PDZ domain binding; Wnt-activated receptor activity; Wnt-activated receptor activity; Wnt-protein binding; protein binding; protein kinase binding; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA