

## Recombinant Mouse FOLH1 cell lysate

**Cat. No.** FOLH1-2452MCL    **Lot. No.** (See product label)

### SPECIFICATION

|                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Product Overview</b>        | Mouse FOLH1 / GCPII / PSMA / NAALAD1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls                                                                                                                                                                                                                                                                            |
| <b>Species</b>                 | Mouse                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| <b>Source</b>                  | Human Cells                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Preparation method</b>      | Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| <b>Lysis buffer</b>            | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF                                                                                                                                                                                                                                                                                                               |
| <b>Quality control Testing</b> | 12.5% SDS-PAGE Stained with Coomassie Blue                                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Recommended Usage</b>       | 1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on                                                                                                 |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

|                            |                                                                                                                                                                                                                                                                                                                                                                                              |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                            | SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.                                                                                                                                                                                                                                                                         |
| <b>Stability</b>           | Samples are stable for up to twelve months from date of receipt at -80°C                                                                                                                                                                                                                                                                                                                     |
| <b>Storage Buffer</b>      | 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF                                                                                                                                                                                                                                                                                    |
| <b>Storage Instruction</b> | Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.                                                                                                                                                                                                        |
| <b>GENE INFORMATION</b>    |                                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Gene Name</b>           | <a href="#">Folh1 folate hydrolase [ Mus musculus ]</a>                                                                                                                                                                                                                                                                                                                                      |
| <b>Official Symbol</b>     | FOLH1                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Synonyms</b>            | FOLH1; folate hydrolase; glutamate carboxypeptidase 2; FGCP; mGCP; GCPII; NAALADase I; folate hydrolase 1; glutamate carboxypeptidase II; membrane glutamate carboxypeptidase; folylpoly-gamma-glutamate carboxypeptidase; prostate-specific membrane antigen homolog; pteroylpoly-gamma-glutamate carboxypeptidase; N-acetylated-alpha-linked acidic dipeptidase I; GCP2; mopsm; MGC141397; |
| <b>Gene ID</b>             | <a href="#">53320</a>                                                                                                                                                                                                                                                                                                                                                                        |
| <b>mRNA Refseq</b>         | <a href="#">NM_001159706</a>                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Protein Refseq</b>      | <a href="#">NP_001153178</a>                                                                                                                                                                                                                                                                                                                                                                 |

|          |                                                                                                                                                                                                                                                                                                          |
|----------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pathway  | One Carbon Metabolism, organism-specific biosystem; Vitamin digestion and absorption, organism-specific biosystem; Vitamin digestion and absorption, conserved biosystem; glutamate removal from folates, organism-specific biosystem; glutathione-mediated detoxification, organism-specific biosystem; |
| Function | carboxypeptidase activity; catalytic activity; dipeptidase activity; hydrolase activity; metal ion binding; metallopeptidase activity; peptidase activity;                                                                                                                                               |

Tel: 1-631-559-9269 1-516-512-3133

Email: [info@creative-biomart.com](mailto:info@creative-biomart.com) Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA