

Recombinant Mouse FOLH1 cell lysate

Cat. No. FOLH1-2452MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse FOLH1 / GCPII / PSMA / NAALAD1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE for Western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	Folh1 folate hydrolase [Mus musculus]
Official Symbol	FOLH1
Synonyms	FOLH1; folate hydrolase; glutamate carboxypeptidase 2; FGCP; mGCP; GCPII; NAALADase I; folate hydrolase 1; glutamate carboxypeptidase II; membrane glutamate carboxypeptidase; folylpoly-gamma-glutamate carboxypeptidase; prostate-specific membrane antigen homolog; pteroylpoly-gamma-glutamate carboxypeptidase; N-acetylated-alpha-linked acidic dipeptidase I; GCP2; mopsm; MGC141397;
Gene ID	53320
mRNA Refseq	NM_001159706
Protein Refseq	NP_001153178

Pathway	One Carbon Metabolism, organism-specific biosystem; Vitamin digestion and absorption, organism-specific biosystem; Vitamin digestion and absorption, conserved biosystem; glutamate removal from folates, organism-specific biosystem; glutathione-mediated detoxification, organism-specific biosystem;
Function	carboxypeptidase activity; catalytic activity; dipeptidase activity; hydrolase activity; metal ion binding; metallopeptidase activity; peptidase activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA