Recombinant Mouse PCSK9 cell lysate

Cat. No. PCSK9-2564MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse PCSK9 / NARC1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Pcsk9 proprotein convertase subtilisin/kexin type 9 [Mus musculus]
Official Symbol	PCSK9
Synonyms	PCSK9; proprotein convertase subtilisin/kexin type 9; NARC-1; convertase subtilisin; proprotein convertase 9; proprotein convertase PC9; subtilisin/kexin-like protease PC9; neural apoptosis regulated convertase 1; neural apoptosis-regulated convertase 1; FH3; PC9; Narc1; HCHOLA3; AI415265; AI747682; MGC47409;
Gene ID	100102
mRNA Refseq	NM_153565
Protein Refseq	NP_705793
Pathway	NGF processing, organism-specific biosystem; Signal Transduction, organism-specific biosystem; Signalling by NGF, organism-specific biosystem;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Function	apolipoprotein receptor binding; hydrolase activity; identical protein binding; low-density lipoprotein particle receptor binding; low-density lipoprotein particle receptor binding; peptidase activity; serine-type endopeptidase activity; serine-type endopeptidase activity; serine-type endopeptidase activity; serine-type peptidase activity; very-low-density lipoprotein particle receptor binding;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA