

Recombinant Human ADAM12 cell lysate

Cat. No. ADAM12-2592HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human ADAM12 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

ADAM12 ADAM metallopeptidase domain 12 [Homo sapiens]

Official Symbol

ADAM12

Synonyms

ADAM12; ADAM metallopeptidase domain 12; a disintegrin and metalloproteinase domain 12 (meltrin alpha); disintegrin and metalloproteinase domain-containing protein 12; MCMPMItna; meltrin alpha; MLTN; meltrin-alpha; MCMP; MLTNA;

Gene ID

8038

mRNA Refseq

NM_003474

Protein Refseq

NP_003465

MIM

602714

UniProt ID

O43184

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	10q26
Pathway	Disease, organism-specific biosystem; Notch signaling pathway, organism-specific biosystem; Signal Transduction, organism-specific biosystem; Signaling by EGFR, organism-specific biosystem; Signaling by EGFR in Cancer, organism-specific biosystem; Syndecan-4-mediated signaling events, organism-specific biosystem;
Function	SH3 domain binding; metal ion binding; metalloendopeptidase activity; metallopeptidase activity; peptidase activity; protein binding; zinc ion binding;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA