

Recombinant Human CD40 cell lysate

Cat. No. CD40-2617HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CD40 / TNFRSF5 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

CD40 CD40 molecule, TNF receptor superfamily member 5 [Homo sapiens]

Official Symbol

CD40

Synonyms

CD40; CD40 molecule, TNF receptor superfamily member 5; TNFRSF5, tumor necrosis factor receptor superfamily, member 5; tumor necrosis factor receptor superfamily member 5; Bp50; p50; CD40L receptor; CD40 type II isoform; B cell-associated molecule; B cell surface antigen CD40; B-cell surface antigen CD40; CD40 antigen (TNF receptor superfamily member 5); tumor necrosis factor receptor superfamily, member 5; nerve growth factor receptor-related B-lymphocyte activation molecule; CDW40; TNFRSF5; MGC9013;

Gene ID

958

mRNA Refseq

NM_001250

Protein Refseq

NP_001241

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	109535
UniProt ID	P25942
Chromosome Location	20q12-q13.2
Pathway	Adaptive Immune System, organism-specific biosystem; Allograft rejection, organism-specific biosystem; Allograft rejection, conserved biosystem; Asthma, organism-specific biosystem; Asthma, conserved biosystem; Autoimmune thyroid disease, organism-specific biosystem; Autoimmune thyroid disease, conserved biosystem;
Function	enzyme binding; protein binding; receptor activity; signal transducer activity;

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA