Recombinant Human LDLR cell lysate

Cat. No. LDLR-2780HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human LDLR / LDL Receptor derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer.Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boiled for 2-5 min.3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles.Notes:The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	LDLR low density lipoprotein receptor [Homo sapiens]
Official Symbol	LDLR
Synonyms	LDLR; low density lipoprotein receptor; low-density lipoprotein receptor; familial hypercholesterolemia; LDL receptor; low-density lipoprotein receptor class A domain-containing protein 3; FH; FHC; LDLCQ2;
Gene ID	3949
mRNA Refseq	NM_000527
Protein Refseq	NP_000518
MIM	606945
UniProt ID	P01130

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	19p13.2
Pathway	Bile secretion, organism-specific biosystem; Bile secretion, conserved biosystem; Chylomicron-mediated lipid transport, organism-specific biosystem; DNA damage response (only ATM dependent), organism-specific biosystem; Endocytosis, organism-specific biosystem; Endocytosis, conserved biosystem; Hepatitis C, organism-specific biosystem;
Function	calcium ion binding; glycoprotein binding; low-density lipoprotein receptor activity; protein binding; receptor activity; very-low-density lipoprotein particle receptor activity;

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA