

Recombinant Human IGFBP3 cell lysate

Cat. No. IGFBP3-2927HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human IGFBP3 / IBP3 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	IGFBP3 insulin-like growth factor binding protein 3 [Homo sapiens]
Official Symbol	IGFBP3
Synonyms	IGFBP3; insulin-like growth factor binding protein 3; insulin-like growth factor-binding protein 3; acid stable subunit of the 140 K IGF complex; binding protein 29; binding protein 53; BP 53; growth hormone dependent binding protein; IBP3; IGF binding protein 3; IBP-3; IGFBP-3; IGF-binding protein 3; growth hormone-dependent binding protein; BP-53;
Gene ID	3486
mRNA Refseq	NM_000598
Protein Refseq	NP_000589
MIM	146732

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P17936
Chromosome Location	7p13-p12
Pathway	Diabetes pathways, organism-specific biosystem; Direct p53 effectors, organism-specific biosystem; Disease, organism-specific biosystem; Myometrial Relaxation and Contraction Pathways, organism-specific biosystem; Regulation of Insulin-like Growth Factor (IGF) Activity by Insulin-like Growth Factor Binding Proteins (IGFBPs), organism-specific biosystem; Transcriptional misregulation in cancer, organism-specific biosystem; Transcriptional misregulation in cancer, conserved biosystem;
Function	insulin-like growth factor I binding; insulin-like growth factor binding; metal ion binding; protein binding; protein tyrosine phosphatase activator activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA