

Recombinant Human DPP4 cell lysate

Cat. No. DPP4-2979HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human DPP4 / CD26 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Human
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.
GENE INFORMATION	
Gene Name	DPP4 dipeptidyl-peptidase 4 [Homo sapiens]
Official Symbol	DPP4
Synonyms	DPP4; dipeptidyl-peptidase 4; ADCP2, adenosine deaminase complexing protein 2 , CD26, dipeptidylpeptidase IV (CD26, adenosine deaminase complexing protein 2); dipeptidyl peptidase 4; DPPIV; ADCP-2; DPP IV; dipeptidylpeptidase 4; dipeptidyl peptidase IV; T-cell activation antigen CD26; adenosine deaminase complexing protein 2; dipeptidylpeptidase IV (CD26, adenosine deaminase complexing protein 2); CD26; ADABP; ADCP2; TP103;
Gene ID	1803
mRNA Refseq	NM_001935
Protein Refseq	NP_001926
MIM	102720

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P27487
Chromosome Location	2q23-qter
Pathway	Incretin Synthesis, Secretion, and Inactivation, organism-specific biosystem; Integration of energy metabolism, organism-specific biosystem; Metabolism, organism-specific biosystem; Protein digestion and absorption, organism-specific biosystem; Protein digestion and absorption, conserved biosystem; Regulation of Insulin Secretion, organism-specific biosystem; Synthesis, Secretion, and Inactivation of Glucagon-like Peptide-1 (GLP-1), organism-specific biosystem;
Function	aminopeptidase activity; collagen binding; dipeptidyl-peptidase activity; peptidase activity; peptide binding; protease binding; protein binding; protein homodimerization activity; receptor activity; receptor binding; serine-type endopeptidase activity; serine-type peptidase activity;