

Recombinant Rat ACVRL1 cell lysate

Cat. No. ACVRL1-3001RCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Rat ALK-1 / ACVRL1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Rat
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reducing agent prior to heating.

Stability

Samples are stable for up to twelve months from date of receipt at -80°C

Storage Buffer

50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF

Storage Instruction

Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name

[Acvrl1](#) activin A receptor type II-like 1 [*Rattus norvegicus*]

Official Symbol

ACVRL1

Synonyms

ACVRL1; activin A receptor type II-like 1; serine/threonine-protein kinase receptor R3; SKR3; TSR-I; Activin receptor likekinase 1; TGF-B superfamily receptor type I; R3; R-3; SETHKIR;

Gene ID

[25237](#)

mRNA Refseq

[NM_022441](#)

Protein Refseq

[NP_071886](#)

Pathway

Id Signaling Pathway, organism-specific biosystem; TGF-beta Receptor Signaling Pathway, organism-specific biosystem;

Function

ATP binding; ATP binding; SMAD binding; SMAD binding; SMAD binding; activin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

binding; activin binding; activin receptor activity, type I; activin receptor activity, type I;
metal ion binding; nucleotide binding; protein serine/threonine kinase activity; receptor
activity; receptor signaling protein serine/threonine kinase activity; transferase activity,
transferring phosphorus-containing groups; transforming growth factor beta binding;
transforming growth factor beta binding; transforming growth factor beta binding;
transforming growth factor beta receptor activity, type I; transforming growth factor
beta receptor activity, type I; contributes_to transforming growth factor beta receptor
activity, type I; transforming growth factor beta-activated receptor activity;
transmembrane receptor protein serine/threonine kinase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA