

Recombinant Mouse ACVRL1 cell lysate

Cat. No. ACVRL1-3093MCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Mouse ALK-1 derived in Human Cells. The whole cell lysate is provided in 1X Sample Buffer. Browse all transfected cell lysate positive controls
Species	Mouse
Source	Human Cells
Preparation method	Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation and then centrifuged to clarify the lysate. The cell lysate was boiled for 5 minutes in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Quality control Testing	12.5% SDS-PAGE Stained with Coomassie Blue
Recommended Usage	1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube. 2. Re-dissolve the pellet using 200µL pure water and boiled for 2-5 min. 3. Store it at -80°C. Recommend to aliquot the cell lysate into smaller quantities for optimal storage. Avoid repeated freeze-thaw cycles. Notes: The lysate is ready to load on SDS-PAGE for Western blot application. If dissociating conditions are required, add

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	reducing agent prior to heating.
Stability	Samples are stable for up to twelve months from date of receipt at -80°C
Storage Buffer	50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF
Storage Instruction	Lysate samples are stable for 12 months from date of receipt when stored at -80°C. Avoid repeated freeze-thaw cycles. Prior to SDS-PAGE fractionation, boil the lysate for 5 minutes.

GENE INFORMATION

Gene Name	Acvrl1 activin A receptor, type II-like 1 [Mus musculus]
Official Symbol	ACVRL1
Synonyms	ACVRL1; activin A receptor, type II-like 1; serine/threonine-protein kinase receptor R3; SKR3; TSR-I; activin receptor-like kinase 1; activin receptor-like kinase-1; TGF-B superfamily receptor type I; Alk1; Acvrlk1; AI115505; AI427544;
Gene ID	11482
mRNA Refseq	NM_009612
Protein Refseq	NP_033742
Pathway	Id Signaling Pathway, organism-specific biosystem; TGF-beta Receptor Signaling Pathway, organism-specific biosystem;
Function	ATP binding; SMAD binding; activin binding; activin receptor activity, type I; kinase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

activity; metal ion binding; nucleotide binding; protein binding; protein kinase activity; protein serine/threonine kinase activity; receptor activity; receptor signaling protein serine/threonine kinase activity; transferase activity; transferase activity, transferring phosphorus-containing groups; transforming growth factor beta binding; transforming growth factor beta receptor activity, type I; transforming growth factor beta-activated receptor activity; transmembrane receptor protein serine/threonine kinase activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA