

Active Recombinant Cynomolgus TNFSF4, Fc-tagged

Cat. No. TNFSF4-85C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	A DNA sequence encoding the cynomolgus TNFSF4 (Gln51-Leu183) was expressed with the Fc region of mouse IgG1 at the N-terminus.
Species	Cynomolgus
Source	Human Cells
ProteinLength	51-183 a.a.
Description	OX-40L, also known as TNFSF4 and CD252, is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. OX-40L is an important costimulatory molecule that plays a crucial role in the regulation of T-cell-mediated immunity. The interaction of TNFSF4-TNFSF4 is involved in the pathogenesis of multiple autoimmune and inflammatory diseases such as systemic lupus erythematosus (SLE), carotid artery disease and cancer. OX-40L is a ligand for receptor TNFRSF4/OX4. It is found to play a role in T cell antigen-presenting cell (APC) interactions. In surface Ig- and CD40-stimulated B cells, this cytokine along with CD70 has been shown to provide CD28-independent costimulatory signals to T cells. This protein and its receptor are reported to directly mediate adhesion of activated T cells to vascular endothelial cells.
Predicted N Terminal	Asp
Form	Lyophilized from sterile PBS, pH 7.4.
Bio-activity	Immobilized Cynomolgus mFc-TNFSF4 at 10 µg/ml (100 µl/well) can bind human

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	TNFRSF4-Fch, The EC50 of human TNFRSF4-Fch is 0.23-0.55 µg/ml.
Molecular Mass	The recombinant cynomolgus TNFSF4 is a disulfide-linked homodimer. The reduced monomer comprises 369 amino acids and has a calculated molecular mass of 42 KDa. The apparent molecular mass of it is approximately 48 KDa respectively in SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method
Purity	> 85 % as determined by SDS-PAGE
Stability	Samples are stable for up to twelve months from date of receipt at -70°C
Storage	Store it under sterile conditions at -70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products.

GENE INFORMATION

Gene Name	TNFSF4 tumor necrosis factor (ligand) superfamily, member 4 [<i>Macaca fascicularis</i> (crab-eating macaque)]
Official Symbol	TNFSF4
Synonyms	TNFSF4; tumor necrosis factor (ligand) superfamily, member 4; hypothetical protein
Gene ID	102141877
mRNA Refseq	XM_005540022

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq	XP_005540079
Chromosome Location	1
Pathway	Cytokine-cytokine receptor interaction