

Active Recombinant Rabbit FCGRT&B2M protein, His-Avi-tagged, Biotinylated

Cat. No. FCGRT&B2M-1419R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Rabbit FCGRT&B2M protein(Ala24-Val297(FCGRT)&Val22-Tyr120(B2M)), fused with His and Avi tag, was expressed in HEK293.
Species	Rabbit
Source	HEK293
ProteinLength	24-297;22-120 a.a.
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4.
Bio-activity	Biotinylated Rabbit FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free captured on Biotin CAP-Series S Sensor Chip can bind Herceptin with an affinity constant of 1.05 µM as determined in a SPR assay (Biacore 8K).Immobilized Biotinylated Rabbit FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free on SA Chip can bind Herceptin with an affinity constant of 1.53 µM as determined in a SPR assay (Biacore 8K).
Molecular Mass	Biotinylated Rabbit FCGRT&B2M Heterodimer Protein, His,Avitag&Tag Free, produced by co-expression of FCGRT and B2M, has a calculated MW of 33.8 kDa (FCGRT) & 11.7 kDa (B2M). Subunit FCGRT is fused with a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™) and subunit Beta-2 microglobulin (B2M)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	contains no tag at the C-terminus. The reducing (R) protein migrates as 35-40 kDa (FCGRT) & 12 kDa (B2M) respectively due to glycosylation.
Proteinlength	Ala24-Val297(FCGRT)&Val22-Tyr120(B2M)
Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>95% as determined by SDS-PAGE. >90% as determined by SEC-MALS.
Storage	For long term, the product should be stored at lyophilized state at -20°C or lower. Please avoid repeated freeze-thaw cycles. This product is stable after at: -20°C to -70°C for 12 months in lyophilized state; -70°C for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin