

Recombinant Human Interleukin 1 Receptor, Type II

Cat. No. IL1R2-1641H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Sf21 Cells
Description	Interleukin 1 receptor type II (IL1R2), also known as CD121b (Cluster of Differentiation 121b), is a transmembrane protein found on B lymphocytes, neutrophils, monocytes, large granular leukocytes and endothelial cells. It is a decoy receptor known for its ability to buffer IL-1 effects. IL-1RII is involved in the control of the proinflammatory state that takes place in the endometrium during the premenstrual and menstrual periods. IL-1RII can neutralize IL-1 beta and counteract its effect on endometrial stromal cells, and may provide a new clinical strategy for the treatment of endometriosis.
State Of Matter	Lyophilized.
Purity	>97% by SDS Page and analyzed by silver stain.
Endotoxin	<1.0 EU/g as determined by the LAL method.
Molecular Weight	The predicted molecular weight of Recombinant Human IL-1 sRII is Mr 36 kDa. However, the actual molecular weight as observed by migration on SDS Page is Mr 40 kDa.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage And Stability

This lyophilized protein is stable for six to twelve months when stored desiccated at -20°C to -70°C. After aseptic reconstitution, this protein may be stored at 2°C to 8°C for one month or at -20°C to -70°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles.

GENE INFORMATION

Gene Name	IL1R2 interleukin 1 receptor, type II [Homo sapiens]
Synonyms	IL1R2; interleukin 1 receptor, type II; IL1RB; CD121b; MGC47725; IL-1R-beta; antigen CDw121b; type II interleukin-1 receptor, beta; Interleukin-1 receptor type II; IL-1R-2; IL-1R-beta; CD121 antigen-like family member B; CD121b antigen; CD121 antigen; interleukin 1 receptor, type II
Gene ID	7850
mRNA Refseq	NM_004633
Protein Refseq	NP_004624
MIM	147811
UniProt ID	P27930
Chromosome Location	2q12
Pathway	Cytokine-cytokine receptor interaction; Hematopoietic cell lineage; MAPK signaling pathway
Function	interleukin-1, Type II, blocking receptor activity; receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA