

Active Recombinant Human IL8 protein, His-Avi-tagged, Biotinylated

Cat. No. IL8-0206H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant Human IL8 protein(Ser28-Ser99), fused with His and Avi tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	28-99 a.a.
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.
Bio-activity	Immobilized Biotinylated Human IL-8, His,Avitag at 1 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate can bind Monoclonal Anti-Human IL-8 Antibody, Human IgG1 with a linear range of 0.06-2 ng/mL.Immobilized Monoclonal Anti-Human IL-8 Antibody, Human IgG1 at 1 µg/mL (100 µL/well) can bind Biotinylated Human IL-8, His,Avitag with a linear range of 0.06-0.2 ng/mL.
Molecular Mass	This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag).The protein has a calculated MW of 12.0 kDa. The protein migrates as 14 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Proteinlength	Ser28-Ser99

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1.0 EU per µg by the LAL method.
Purity	>90% as determined by SDS-PAGE.
Storage	For long term, the product should be stored at lyophilized state at -20°C or lower. Please avoid repeated freeze-thaw cycles. This product is stable after at: -20°C to -70°C for 12 months in lyophilized state; -70°C for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.
Conjugation	Biotin

GENE INFORMATION

Gene Name	IL8 interleukin 8 [Homo sapiens]
Official Symbol	IL8
Synonyms	IL8; interleukin 8; interleukin-8; 3 10C; alveolar macrophage chemotactic factor I; AMCF I; b ENAP; beta endothelial cell derived neutrophil activating peptide; chemokine (C X C motif) ligand 8; CXCL8; GCP 1; GCP1; granulocyte chemotactic protein 1; IL 8; K60; LECT; LUCT; lung giant cell carcinoma derived chemotactic protein; lymphocyte derived neutrophil activating peptide; LYNAP; MDNCF; MONAP; monocyte derived neutrophil chemotactic factor; monocyte derived neutrophil activating peptide; NAF; NAP 1; NAP1; neutrophil activating peptide 1; SCYB8; TSG 1; tumor necrosis factor induced gene 1; emoctakin; T-cell chemotactic factor; neutrophil-activating peptide 1; chemokine (C-X-C motif) ligand 8; beta-thromboglobulin-like protein; tumor necrosis factor-induced gene 1; monocyte-derived neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; small

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

inducible cytokine subfamily B, member 8; lung giant cell carcinoma-derived chemotactic protein; beta endothelial cell-derived neutrophil activating peptide; GCP-1; NAP-1;

Gene ID [3576](#)

mRNA Refseq [NM_000584](#)

Protein Refseq [NP_000575](#)

MIM [146930](#)

UniProt ID [P10145](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA