Rhesus monkey Ileum Lysate

Cat. No. Ileum-245R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Rhesus macaque
Source	Rhesus Small Intestine: Ileum
Concentration	2 mg/mL or better
Tissue Type	Small intestine: Ileum
Shipping	Dry ice
Preparation method	Monkey (Rhesus) Ileum tissue lysate was prepared by homogenization using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70?C. The monkey (Rhesus) Ileum tissue total protein is provided in a buffer including HEPES (pH 7.9), MgCl2, KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, NP-40, and a cocktail of protease inhibitors. For quality control purposes, the Ileum tissue pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The Ileum tissue is then Western analyzed by either GAPDH or β-actin antibody, and the expression level is consistent with each lot.
Diagnosis	Normal
Storage Buffer	HEPES pH 7.9, MgCl2, KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	NP-40.
Storage Instruction	Store at 2-8?C for continuous use. For extended storage, freeze working aliquots at -70?C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.
Applications	Ileum lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA