

Human Lymphoma, Hodgkins Disease Membrane Tumor Lysate

Cat. No. Lymphoma-334H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Human lymphoma, Hodgkins Disease
Concentration	2 mg/mL or better
Tissue Type	lymphoma, Hodgkins Disease
Shipping	Dry ice
Preparation method	Human lymphoma, Hodgkins Disease tissue membrane protein lysate was prepared by isolating the membrane protein from whole tissue homogenates using a proprietary technique. The human lymphoma, Hodgkins Disease tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. The membrane protein is provided in a buffer including HEPES (pH 7.9), MgCl₂, KCl, EDTA, Sucrose, Glycerol, sodium deoxycholate, NP-40, and a cocktail of protease inhibitors. For quality control purposes, the isolated lymphoma, Hodgkins Disease tissue membrane protein pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The isolated lymphoma, Hodgkins Disease tissue membrane protein is then Western analyzed by either GAPDH or β-actin antibody to confirm there is no signal or very weak signal.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Diagnosis	Tumor
Storage Buffer	HEPES pH 7.9, MgCl ₂ , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and NP-40.
Storage Instruction	Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.
Applications	lymphoma, Hodgkins Disease membrane protein lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA