

# Rhesus monkey Occipital Lobe Lysate

**Cat. No.** Occipital lobe-348R    **Lot. No.** (See product label)

## SPECIFICATION

### Product Overview

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species</b>            | Rhesus macaque                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Source</b>             | Rhesus Brain: Occipital Lobe                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Concentration</b>      | 2 mg/mL or better                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <b>Tissue Type</b>        | Brain: Occipital Lobe                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>Shipping</b>           | Dry ice                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Preparation method</b> | <p>Monkey (Rhesus) occipital lobe tissue lysate was prepared by homogenization using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. The monkey (Rhesus) occipital lobe tissue total protein is provided in a buffer including HEPES (pH 7.9), MgCl<sub>2</sub>, KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, NP-40, and a cocktail of protease inhibitors. For quality control purposes, the occipital lobe tissue pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The occipital lobe tissue is then Western analyzed by either GAPDH or <math>\beta</math>-actin antibody, and the expression level is consistent with each lot.</p> |
| <b>Diagnosis</b>          | Normal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Storage Buffer</b>     | HEPES pH 7.9, MgCl <sub>2</sub> , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |

 Tel: 1-631-559-9269    1-516-512-3133

 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)     Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

NP-40.

**Storage Instruction**

Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.

**Applications**

Occipital Lobe lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133 Email: [info@creative-biomart.com](mailto:info@creative-biomart.com)  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA