

CD105 protein-coupled magnetic MicroBeads

Cat. No. CD105-403M Lot. No. (See product label)

SPECIFICATION

Product Overview

Species Human

Capacity for 1×10^9 total cells

Background

The CD105 antigen (endoglin) serves as a receptor for the growth and differentiation factors TGF- β 1 and TGF- β 3, and is expressed on mature endothelial cells, on MSCs with mesodermal differentiation capacity, and on some leukemic cells of B lymphoid and myeloid origin. The CD105 MicroBeads recognize the same molecular antigen as anti-SH2 antibodies-the most commonly employed antigen for the definition of cultivated MSCs.

Application


CD105 MicroBeads can be used for studies on mesengensis and for in vitro investigations on hematological disorders. CD105+ bone marrow cells also show multipotent differentiation in vitro and the capacity to form bone in vivo without prior cultivation or differentiation.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Analysis



Separation of CD105+ endothelial cells from a single-cell preparation of umbilical cord vein tissue using CD105 MicroBeads, an MS Column, and a MiniMACS Separator.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA