

SiglecF Antibody-coupled magnetic MicroBeads

Cat. No. SiglecF-503M **Lot. No.** (See product label)

SPECIFICATION

Product Overview


Species Mouse

Capacity for 1–2×10⁹ cells

Background

Siglec-F, also known as Siglec-5, is a member of the sialic acid-binding immunoglobulin (Ig)-like lectins (Siglecs). The extracellular Ig domain binds specifically carbohydrate structures containing sialic acid residues. Siglec-F has, like most Siglecs, an immunoreceptor tyrosine-based inhibitory motif (ITIM) in its cytoplasmic domain and is discussed to play a role in eosinophil accumulation and survival. It is expressed in a tissue dependent manner on most eosinophils, but also found on alveolar macrophages and some neutrophils.

Analysis



Siglec-F⁺ eosinophils were isolated from mouse peripheral blood using Anti-Siglec-F MicroBeads, two MS Columns, and a MiniMACS Separator. Cells were fluorescently stained with Anti-Siglec-F-APC and Anti-Gr-1-VioGreen and analyzed by flow cytometry using the MACSQuant Analyzer. Cell debris and dead cells were excluded from the analysis based on scatter signals and propidium iodide

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

fluorescence.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA