

Human Prostate Liver Cirrhosis Lysate

Cat. No. Prostate Liver Cirrhosis -403H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Human Prostate
Concentration	2 mg/mL or better
Tissue Type	Prostate
Shipping	Dry ice
Preparation method	Human prostate tissue lysate was prepared by homogenization using a proprietary technique. The tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. The human prostate tissue total protein is provided in a buffer including HEPES (pH7.9), MgCl ₂ , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, NP-40, and a cocktail of protease inhibitors. For quality control purposes, the prostate tissue pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The prostate tissue is then Western analyzed by either GAPDH or β -actin antibody, and the expression level is consistent with each lot.
Diagnosis	Diseased
Storage Buffer	HEPES pH 7.9, MgCl ₂ , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

NP-40.

Storage Instruction

Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.

Applications

Prostate lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA