Human Raji Cytoplasmic Lysate

Cat. No. Raji-407H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Species	Human
Source	Raji Cell Line
Concentration	2 mg/mL or better
Cell line	Raji (Human Lymphoma; B Lymphoma)
Shipping	Dry ice
Preparation method	Raji (Human lymphoma; B lymphoma) Cell cytoplasmic protein lysate cytoplasmic protein lysate was prepared by isolating the cytoplasmic protein from whole tissue homogenates using a proprietary technique. The cytoplasmic protein is provided in a buffer including HEPES (pH 7.9), MgCl2, KCl, EDTA, Sucrose, glycerol, and a cocktail of protease inhibitors. For quality control purposes, the isolated Raji (Human lymphoma; B lymphoma) cytoplasmic protein pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The isolated Raji (Human lymphoma; B lymphoma) tissue cytoplasmic protein is then Western analyzed by GAPDH antibody, and the expression level is consistent with each lot.
Diagnosis	Tumor
Storage Buffer	HEPES pH 7.9, MgCl2, KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

	NP-40.
Storage Instruction	Store at 2-8?C for continuous use. For extended storage, freeze working aliquots at -70?C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.
Applications	Raji (Lymphoma; B Lymphoma) Cell Cytoplasmic protein lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA