

Human Spleen Cytoplasmic Lysate

Cat. No. Spleen-465H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	Human Spleen
Concentration	2 mg/mL or better
Tissue Type	Spleen
Shipping	Dry ice
Preparation method	Human spleen tissue cytoplasmic protein lysate was prepared by isolating the cytoplasmic protein from whole tissue homogenates using a proprietary technique. The human spleen tissue was frozen in liquid nitrogen immediately after excision and then stored at -70°C. The cytoplasmic protein is provided in a buffer including HEPES (pH 7.9), MgCl ₂ , KCl, EDTA, Sucrose, glycerol, and a cocktail of protease inhibitors. For quality control purposes, the isolated spleen tissue cytoplasmic protein pattern on SDS-PAGE gel is shown to be consistent for each lot by visualization with coomassie blue staining. The isolated spleen tissue cytoplasmic protein is then Western analyzed by GAPDH antibody, and the expression level is consistent with each lot.
Diagnosis	Normal
Storage Buffer	HEPES pH 7.9, MgCl ₂ , KCl, EDTA, Sucrose, Glycerol, Sodium deoxycholate, and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

NP-40.

Storage Instruction

Store at 2-8°C for continuous use. For extended storage, freeze working aliquots at -70°C. Repeated freezing and thawing is not recommended. Under proper storage conditions the shelf life is half a year from the date of receipt.

Applications

Spleen cytoplasmic protein lysate is for use in Western blotting, 10 µg to 20 µg per lane is recommended for mini gel.

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA