

Leukopak CD4 protein-coupled magnetic MicroBeads

Cat. No. CD4-3913M **Lot. No.** (See product label)

SPECIFICATION

Product Overview


Species Human

Capacity for 0.5 Leukopaks

Background The CD4 antigen is expressed on T helper cells and at a lower level on monocytes and dendritic cells. The CD4 antigen is an accessory molecule in the recognition of MHC class II/peptide complexes by the TCR heterodimer on CD4+ T helper cells.

Application The purified CD4+ T cells are well suited for further flow cytometric, functional, or molecular analysis.

Analysis



Separation of a Leukopak sample using the StraightFrom Leukopak CD4 MicroBead Kit and the MultiMACS Cell24 Separator Plus with the Multi-24 Column Blocks. Cells were fluorescently stained with CD4-PE, CD14-APC, as well as CD45-VioBlue and analyzed by flow cytometry using the MACSQuant Analyzer. Cell debris and dead cells were excluded from the analysis based on scatter signals and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

propidium iodide fluorescence.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA