

Active Recombinant Bovine FGF2 protein, His-tagged

Cat. No. FGF2-019B **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Bovine FGF2 protein(Pro10~Ser155), fused with N-terminal His Tag, was expressed in E. coli.
Species	Bovine
Source	E.coli
ProteinLength	Pro10~Ser155
Form	20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.
Bio-activity	Basic fibroblast growth factor (FGF2), also known as bFGF, FGF-β is a member of a large family of structurally related heparin-binding proteins (the FGFs) involved in the regulation of cell proliferation, growth and differentiation. It involved in many biological processes including angiogenesis, embryonic development and wound healing. Additionally, FGF2 is a critical component of human embryonic stem cell culture medium. Besides, Caspase 1 (CASP1) has been identified as an interactor of FGF2, thus a binding ELISA assay was conducted to detect the interaction of recombinant bovine FGF2 and recombinant bovine CASP1. Briefly, FGF2 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to CASP1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-FGF2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

minutes at 37°C. Finally, add 50μ stop solution to the wells and read at 450nm immediately. The binding activity of FGF2 and CASP1 was shown in Figure 1, and this effect was in a dose dependent manner.


Molecular Mass 20.1kDa

Purity > 97%

Storage Avoid repeated freeze/thaw cycles. Store at 2-8°C for one month. Aliquot and store at -80°C for 12 months.

Reconstitution Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.


Activity



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA