

Recombinant Human CYP2A6, MYC/DDK-tagged

Cat. No. CYP2A6-126H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CYP2A6, fused with C-terminal MYC/DDK, was expressed in HEK293 cells.
Species	Human
Source	HEK293
Description	This gene, CYP2A6, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by phenobarbital. The enzyme is known to hydroxylate coumarin, and also metabolizes nicotine, aflatoxin B1, nitrosamines, and some pharmaceuticals. Individuals with certain allelic variants are said to have a poor metabolizer phenotype, meaning they do not efficiently metabolize coumarin or nicotine. This gene is part of a large cluster of cytochrome P450 genes from the CYP2A, CYP2B and CYP2F subfamilies on chromosome 19q. The gene was formerly referred to as CYP2A3; however, it has been renamed CYP2A6.
Molecular Mass	56.3 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Buffer 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10% glycerol.

GENE INFORMATION

Gene Name CYP2A6 cytochrome P450, family 2, subfamily A, polypeptide 6 [Homo sapiens]

Official Symbol CYP2A6

Synonyms CYP2A6; CPA6; CYP2A; CYP2A3; P450PB; CYP2A6; P450C2A; cytochrome P450, family 2, subfamily A, polypeptide 6; cytochrome P450 2A6; cytochrome P450(I); cytochrome P450 IIA3; coumarin 7-hydroxylase; xenobiotic monooxygenase; 1,4-cineole 2-exo-monooxygenase; flavoprotein-linked monooxygenase; cytochrome P450, subfamily IIA (phenobarbital-inducible), polypeptide 6; EC 1.14.14.1

Gene ID 1548

mRNA Refseq NM_000762

Protein Refseq NP_000753

MIM 122720

UniProt ID P11509

Chromosome Location 19q13.2

Pathway Biological oxidations; Caffeine metabolism; Chemical carcinogenesis

Function coumarin 7-hydroxylase activity; oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen, reduced flavin or flavoprotein as one

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

donor, and incorporation of one atom of oxygen; enzyme binding