

Recombinant Human XRCC5 293 Cell Lysate

Cat. No. XRCC5-254HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining) (XRCC5) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name XRCC5 X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining) [Homo sapiens]

Official Symbol XRCC5

Synonyms XRCC5; X-ray repair complementing defective repair in Chinese hamster cells 5 (double-strand-break rejoining); X ray repair complementing defective repair in Chinese hamster cells 5 (double strand break rejoining; Ku autoantigen, 80kD); X-ray repair cross-complementing protein 5; KARP 1; Ku autoantigen; 80kDa; KU80; Ku86; KUB2; TLAA; CTC85; CTCBF; nuclear factor IV; Ku autoantigen, 80kDa; DNA repair protein XRCC5; thyroid-lupus autoantigen; 86 kDa subunit of Ku antigen; lupus Ku autoantigen protein p86; Ku86 autoantigen related protein 1; CTC box-binding factor 85 kDa subunit; ATP-dependent DNA helicase 2 subunit 2; ATP-dependent DNA helicase II 80 kDa subunit; NFIV; KARP1; KARP-1; FLJ39089;

Gene ID 7520

mRNA Refseq NM_021141

Protein Refseq NP_066964

MIM 194364

UniProt ID P13010

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	2q35
Pathway	2-LTR circle formation, organism-specific biosystem; BARD1 signaling events, organism-specific biosystem; Coregulation of Androgen receptor activity, organism-specific biosystem; DNA Repair, organism-specific biosystem; DNA-PK complex, organism-specific biosystem; Disease, organism-specific biosystem; Double-Strand Break Repair, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA