

Recombinant Human FCGR3A Protein, V176F, Gly17-Gln208, C-6×His tagged

Cat. No. FCGR3A-045H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human FCGR3A Protein, Gly17-Gln208, with a C-terminal 6-His tag was expressed in Human embryonic kidney cell.
Species	Human
Source	HEK293
ProteinLength	Gly17-Gln208
Description	This gene encodes a receptor for the Fc portion of immunoglobulin G, and it is involved in the removal of antigen-antibody complexes from the circulation, as well as other responses, including antibody dependent cellular mediated cytotoxicity and antibody dependent enhancement of virus infections. This gene (FCGR3A) is highly similar to another nearby gene (FCGR3B) located on chromosome 1. The receptor encoded by this gene is expressed on natural killer (NK) cells as an integral membrane glycoprotein anchored through a transmembrane peptide, whereas FCGR3B is expressed on polymorphonuclear neutrophils (PMN) where the receptor is anchored through a phosphatidylinositol (PI) linkage. Mutations in this gene are associated with immunodeficiency 20, and have been linked to susceptibility to recurrent viral infections, susceptibility to systemic lupus erythematosus, and alloimmune neonatal neutropenia. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bio-activity	Measured by its binding ability in a functional ELISA. When Recombinant Human Fc gamma RIIIa/CD16a (V176F) was immobilize on a Ms x PolyHis coated plate, it binds biotinylated Human IgG. The concentration of biotinylated Human IgG that produces 50% of the optimal binding response is approximately 0.5-2.5 µg/mL.
Molecular Mass	Predicted Molecular Mass: 23 kDa SDS-PAGE: 43-58 kDa, reducing conditions
Endotoxin	<0.10 EU/µg of the protein by the LAL method.
Purity	>95%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie Blue Staining.
N-terminal Sequence Analysis	Gly17 & Met18
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from a 0.2 µm filtered solution in sterile PBS.
Shipping	The product is shipped at ambient temperature.
Reconstitution	Reconstitute at 100 µg/mL in PBS.
GENE INFORMATION	
Gene Name	FCGR3A Fc gamma receptor IIIa [Homo sapiens (human)]

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	FCGR3A
Synonyms	FCGR3A; Fc gamma receptor IIIa; CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRIII; CD16-II; FCGRIII; FCRIIIA; FcGRIIIA; low affinity immunoglobulin gamma Fc region receptor III-A; CD16a antigen; Fc fragment of IgG receptor IIIa; Fc fragment of IgG, low affinity III, receptor for (CD16); Fc fragment of IgG, low affinity IIIa, receptor (CD16a); Fc gamma receptor III-A; Fc-gamma RIII-alpha; Fc-gamma receptor III-2 (CD 16); Fc-gamma receptor IIIb (CD16); FcgammaRIIIA; IgG Fc receptor III-A; igG Fc receptor III-2; immunoglobulin G Fc receptor III; low affinity immunoglobulin gamma Fc region receptor III-A/III-B recombinant; low affinity immunoglobulin gamma Fc region receptor III-B; low affinity immunoglobulin gamma receptor III-a Fc fragment; neutrophil-specific antigen NA
Gene ID	2214
mRNA Refseq	NM_000569
Protein Refseq	NP_000560
MIM	146740
UniProt ID	P08637

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA