

Recombinant Human VKORC1 293 Cell Lysate

Cat. No. VKORC1-405HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source HEK293

Description

Antigen standard for vitamin K epoxide reductase complex, subunit 1 (VKORC1), transcript variant 1 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.

Components

This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).

Size 0.1 mg

Storage Instruction

Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION
Gene Name

VKORC1 vitamin K epoxide reductase complex, subunit 1 [Homo sapiens]

Official Symbol

VKORC1

Synonyms

VKORC1; vitamin K epoxide reductase complex, subunit 1; vitamin K dependent clotting factors deficiency 2 , VKCFD2; vitamin K epoxide reductase complex subunit 1; phyloquinone epoxide reductase; vitamin K1 2,3-epoxide reductase subunit 1; vitamin K dependent clotting factors deficiency 2; vitamin K1 epoxide reductase (warfarin-sensitive); VKOR; MST134; MST576; VKCFD2; EDTP308; IMAGE3455200; MGC2694; FLJ00289;

Gene ID

79001

mRNA Refseq

NM_024006

Protein Refseq

NP_076869

MIM

608547

UniProt ID

Q9BQB6

Chromosome Location

16p11.2

Pathway

Gamma-carboxylation, transport, and amino-terminal cleavage of proteins, organism-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

specific biosystem; Metabolism of proteins, organism-specific biosystem; PTM: gamma carboxylation, hypusine formation and arylsulfatase activation, organism-specific biosystem; Post-translational protein modification, organism-specific biosystem;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA