

Active Recombinant Bovine HGF Protein, NK1 isoform, Animal-Free

Cat. No. HGF-03B **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant bovine HGF protein is a potent, high-purity NK1 isoform of bovine hepatocyte growth factor (HGF). Animal-free (AOF) and carrier protein-free.
Species	Bovine
Source	E.coli
Description	Hepatocyte growth factor (HGF) regulates cell growth, cell motility, and morphogenesis and plays a central role in angiogenesis, tumorigenesis, and tissue regeneration. HGF binds and activates the receptor tyrosine kinase, c-Met, activating PI3K/AKT, FAK, JNK, and ERK1/2 signaling.
Form	Lyophilized from acetonitrile, TFA
Molecular Mass	20 kDa
Endotoxin	< 0.005 EU/μg protein (below level of detection)
Purity	> 98%, by SDS-PAGE quantitative densitometry
Applications	Differentiation of iPSCs to hepatocyte-like cells; iPSC and primary cell differentiation to muscle; Cellular agriculture and cultivated meat cell culture media optimization
Reconstitution	Resuspend in 10mM HCl at > 100 μg/mL (provided with protein and free of charge),

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

prepare single use aliquots, add carrier protein if desired and store frozen at -20 or -80 centigrade

GENE INFORMATION

Gene Name HGF hepatocyte growth factor [Bos taurus (cattle)]

Official Symbol HGF

Synonyms HGF; hepatocyte growth factor; hepatocyte growth factor; SF; hepapoietin A; hepatopoeitin-A; hepatopietin-A; scatter factor


Gene ID 282879

mRNA Refseq NM_001031751

Protein Refseq NP_001026921

UniProt ID F1MZD6

Bioactivity of Bovine hepatocyte growth factor




Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

HGF NK1 activity is determined using the Promega serum response element luciferase reporter assay in HEK293T cells. EC₅₀ = 2.0 ng/mL (96 pM). Cells are treated in triplicate with a serial dilution of HGF NK1 for 3 hours. Firefly luciferase activity is measured and normalized to the control Renilla luciferase activity. Please note: protein activity has been determined in a standardized assay using human cells.

Purity



Recombinant bovine HGF protein migrates as a single band at 18 kDa in non-reducing (NR) conditions and 20 kDa upon reduction (R). Purified recombinant protein (3 µg) was resolved using 15% w/v SDS-PAGE in reduced (+β-mercaptoethanol, R) and non-reduced (NR) conditions and stained with Coomassie Brilliant Blue R250.