

Recombinant Human UBA52 293 Cell Lysate

Cat. No. UBA52-603HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for ubiquitin A-52 residue ribosomal protein fusion product 1 (UBA52), transcript variant 2 is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications

ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name [UBA52 ubiquitin A-52 residue ribosomal protein fusion product 1 \[Homo sapiens \]](#)

Official Symbol UBA52

Synonyms

UBA52; ubiquitin A-52 residue ribosomal protein fusion product 1; ubiquitin-60S ribosomal protein L40; 60S ribosomal protein L40; CEP52; HUBCEP52; L40; MGC57125; MGC126879; MGC126881; ribosomal protein L40; RPL40; ubiquitin carboxyl extension protein 52; ubiquitin 52 amino acid fusion protein; ubiquitin CEP52; ubiquitin-CEP52; ubiquitin-52 amino acid fusion protein;

Gene ID [7311](#)

mRNA Refseq [NM_001033930](#)

Protein Refseq [NP_001029102](#)

MIM [191321](#)

UniProt ID [P62987](#)

Chromosome Location 19p13.1-p12

Pathway

APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of Cyclin B, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem;
APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem;
APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted
proteins in late mitosis/early G1, organism-specific biosystem; Activated NOTCH1
Transmits Signal to the Nucleus, organism-specific biosystem; Activated NOTCH1
Transmits Signal to the Nucleus, organism-specific biosystem;