

Recombinant Human TUBB2B 293 Cell Lysate

Cat. No. TUBB2B-650HCL **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Human
Source	HEK293
Description	Antigen standard for tubulin, beta 2B (TUBB2B) is a lysate prepared from HEK293T cells transiently transfected with a TrueORF gene-carrying pCMV plasmid and then lysed in RIPA Buffer. Protein concentration was determined using a colorimetric assay. The antigen control carries a C-terminal Myc/DDK tag for detection.
Components	This product includes 3 vials: 1 vial of gene-specific cell lysate, 1 vial of control vector cell lysate, and 1 vial of loading buffer. Each lysate vial contains 0.1 mg lysate in 0.1 ml (1 mg/ml) of RIPA Buffer (50 mM Tris-HCl pH7.5, 250 mM NaCl, 5 mM EDTA, 50 mM NaF, 1% NP40). The loading buffer vial contains 0.5 ml 2X SDS Loading Buffer (125 mM Tris-Cl, pH6.8, 10% glycerol, 4% SDS, 0.002% Bromophenol blue, 5% beta-mercaptoethanol).
Size	0.1 mg
Storage Instruction	Store at -80°C. Minimize freeze-thaw cycles. After addition of 2X SDS Loading Buffer, the lysates can be stored at -20°C. Product is guaranteed 6 months from the date of shipment.
Applications	ELISA, WB, IP. WB: Mix equal volume of lysates with 2X SDS Loading Buffer. Boil

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

the mixture for 10 min before loading (for membrane protein lysates, incubate the mixture at room temperature for 30 min). Load 5 ug lysate per lane.

GENE INFORMATION

Gene Name	TUBB2B tubulin, beta 2B class IIb [Homo sapiens]
Official Symbol	TUBB2B
Synonyms	TUBB2B; tubulin, beta 2B class IIb; tubulin, beta 2B; tubulin beta-2B chain; bA506K6.1; class IIb beta tubulin; DKFZp566F223; MGC8685; class IIb beta-tubulin; class II beta-tubulin isotype; tubulin, beta polypeptide paralog; RP11-506K6.1; FLJ98847;
Gene ID	347733
mRNA Refseq	NM_178012
Protein Refseq	NP_821080
MIM	612850
UniProt ID	Q9BVA1
Chromosome Location	6p25.2
Pathway	Chaperonin-mediated protein folding, organism-specific biosystem; Cooperation of Prefoldin and TriC/CCTin actin and tubulin folding, organism-specific biosystem; Formation of tubulin folding intermediates by CCT/TriC, organism-specific biosystem; Gap junction, organism-specific biosystem; Gap junction, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Metabolism of proteins, organism-specific biosystem; Pathogenic Escherichia coli infection, organism-specific biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA